



Nicolaides-Baraitser syndrome

Nicolaides-Baraitser syndrome is a condition that affects many body systems. Affected individuals can have a wide variety of signs and symptoms, but the most common are sparse scalp hair, small head size (microcephaly), distinct facial features, short stature, prominent finger joints, unusually short fingers and toes (brachydactyly), recurrent seizures (epilepsy), and moderate to severe intellectual disability with impaired language development.

In people with Nicolaides-Baraitser syndrome, the sparse scalp hair is often noticeable in infancy. The amount of hair decreases over time, but the growth rate and texture of the hair that is present is normal. Affected adults generally have very little hair. In rare cases, the amount of scalp hair increases over time. As affected individuals age, their eyebrows may become less full, but their eyelashes almost always remain normal. At birth, the hair on the face may be abnormally thick (hypertrichosis) but thins out over time.

Most affected individuals grow slowly, resulting in short stature and microcephaly. Sometimes, growth before birth is unusually slow.

The characteristic facial features of people with Nicolaides-Baraitser syndrome include a triangular face, deep-set eyes, a thin nasal bridge, wide nostrils, a pointed nasal tip, and a thick lower lip. Many affected individuals have a lack of fat under the skin (subcutaneous fat) of the face, which may cause premature wrinkling. Throughout their bodies, people with Nicolaides-Baraitser syndrome may have pale skin with veins that are visible on the skin surface due to the lack of subcutaneous fat.

In people with Nicolaides-Baraitser syndrome, a lack of subcutaneous fat in the hands makes the finger joints appear larger than normal. Over time, the fingertips become broad and oval shaped. Additionally, there is a wide gap between the first and second toes (known as a sandal gap).

Most people with Nicolaides-Baraitser syndrome have epilepsy, which often begins in infancy. Affected individuals can experience multiple seizure types, and the seizures can be difficult to control with medication.

Almost everyone with Nicolaides-Baraitser syndrome has moderate to severe intellectual disability. Early developmental milestones, such as crawling and walking, are often normally achieved, but further development is limited, and language development is severely impaired. At least one-third of affected individuals never develop speech, while others lose their verbal communication over time. People with this condition are often described as having a happy demeanor and being very friendly, although they can exhibit moments of aggression and temper tantrums.

Other signs and symptoms of Nicolaides-Baraitser syndrome include an inflammatory skin disorder called eczema. About half of individuals with Nicolaides-Baraitser syndrome have a soft out-pouching around the belly-button (umbilical hernia) or lower abdomen (inguinal hernia). Some affected individuals have dental abnormalities such as widely spaced teeth, delayed eruption of teeth, and absent teeth (hypodontia). Most affected males have undescended testes (cryptorchidism) and females may have underdeveloped breasts. Nearly half of individuals with Nicolaides-Baraitser syndrome have feeding problems.

Frequency

Nicolaides-Baraitser syndrome is likely a rare condition; approximately 75 cases have been reported in the scientific literature.

Genetic Changes

Nicolaides-Baraitser syndrome is caused by mutations in the *SMARCA2* gene. This gene provides instructions for making one piece (subunit) of a group of similar protein complexes known as SWI/SNF complexes. These complexes regulate gene activity (expression) by a process known as chromatin remodeling. Chromatin is the network of DNA and proteins that packages DNA into chromosomes. The structure of chromatin can be changed (remodeled) to alter how tightly DNA is packaged. Chromatin remodeling is one way gene expression is regulated during development; when DNA is tightly packed, gene expression is lower than when DNA is loosely packed. To provide energy for chromatin remodeling, the *SMARCA2* protein uses a molecule called ATP.

The *SMARCA2* gene mutations that cause Nicolaides-Baraitser syndrome result in the production of an altered protein that interferes with the normal function of the SWI/SNF complexes. These altered proteins are able to form SWI/SNF complexes, but the complexes are nonfunctional. As a result, they cannot participate in chromatin remodeling. Disturbance of this regulatory process alters the activity of many genes, which likely explains the diverse signs and symptoms of Nicolaides-Baraitser syndrome.

Inheritance Pattern

Nicolaides-Baraitser syndrome follows an autosomal dominant pattern of inheritance, which means one copy of the altered gene in each cell is sufficient to cause the disorder.

All cases of this condition result from new (de novo) mutations in the gene that occur during the formation of reproductive cells (eggs or sperm) or in early embryonic development. These cases occur in people with no history of the disorder in their family.

Other Names for This Condition

- NBS
- NCBRS

Diagnosis & Management

Genetic Testing

- Genetic Testing Registry: Nicolaides-Baraitser syndrome
<https://www.ncbi.nlm.nih.gov/gtr/conditions/C1303073/>

Other Diagnosis and Management Resources

- GeneReview: Nicolaides-Baraitser Syndrome
<https://www.ncbi.nlm.nih.gov/books/NBK321516>

General Information from MedlinePlus

- Diagnostic Tests
<https://medlineplus.gov/diagnostictests.html>
- Drug Therapy
<https://medlineplus.gov/drugtherapy.html>
- Genetic Counseling
<https://medlineplus.gov/geneticcounseling.html>
- Palliative Care
<https://medlineplus.gov/palliativecare.html>
- Surgery and Rehabilitation
<https://medlineplus.gov/surgeryandrehabilitation.html>

Additional Information & Resources

MedlinePlus

- Encyclopedia: Microcephaly
<https://medlineplus.gov/ency/article/003272.htm>
- Encyclopedia: Short Stature
<https://medlineplus.gov/ency/article/003271.htm>
- Health Topic: Developmental Disabilities
<https://medlineplus.gov/developmentaldisabilities.html>
- Health Topic: Seizures
<https://medlineplus.gov/seizures.html>
- Health Topic: Speech and Language Problems in Children
<https://medlineplus.gov/speechandlanguageproblemsinchildren.html>

Genetic and Rare Diseases Information Center

- Nicolaides-Baraitser syndrome
<https://rarediseases.info.nih.gov/diseases/270/nicolaides-baraitser-syndrome>

Educational Resources

- Centers for Disease Control and Prevention: Developmental Disabilities
<https://www.cdc.gov/ncbddd/developmentaldisabilities/facts.html>
- Disease InfoSearch: Nicolaides Baraitser syndrome
<http://www.diseaseinfosearch.org/Nicolaides+Baraitser+syndrome/5221>
- Kennedy Krieger Institute: Intellectual Disability
<https://www.kennedykrieger.org/patient-care/diagnoses-disorders/intellectual-disability>
- MalaCards: nicolaides-baraitser syndrome
http://www.malacards.org/card/nicolaides_baraitser_syndrome_2
- Merck Manual Consumer Version: Seizure Disorders
<http://www.merckmanuals.com/home/brain-spinal-cord-and-nerve-disorders/seizure-disorders/seizure-disorders>
- Orphanet: Intellectual disability-sparse hair-brachydactyly syndrome
http://www.orpha.net/consor/cgi-bin/OC_Exp.php?Lng=EN&Expert=3051

Patient Support and Advocacy Resources

- American Association on Intellectual and Developmental Disabilities (AAIDD)
<http://aaidd.org/>
- American Epilepsy Society
<https://www.aesnet.org/>
- Citizens United for Research in Epilepsy (CURE)
<http://www.cureepilepsy.org/home.asp>
- North American Hair Research Society
<http://nahrs.org/Home.aspx>
- The Arc
<http://www.thearc.org/>
- The MAGIC Foundation
<https://www.magicfoundation.org/>

GeneReviews

- Nicolaides-Baraitser Syndrome
<https://www.ncbi.nlm.nih.gov/books/NBK321516>

Scientific Articles on PubMed

- PubMed
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28Nicolaides-Baraitser+syndrome%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+3600+days%22%5Bdp%5D>

OMIM

- NICOLAIDES-BARAITSER SYNDROME
<http://omim.org/entry/601358>

Sources for This Summary

- Castori M, Covaciu C, Rinaldi R, Grammatico P, Paradisi M. A rare cause of syndromic hypotrichosis: Nicolaides-Baraitser syndrome. *J Am Acad Dermatol*. 2008 Nov;59(5 Suppl):S92-8. doi: 10.1016/j.jaad.2008.05.016.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/19119135>
- Sousa SB, Abdul-Rahman OA, Bottani A, Cormier-Daire V, Fryer A, Gillesen-Kaesbach G, Horn D, Josifova D, Kuechler A, Lees M, MacDermot K, Magee A, Morice-Picard F, Rosser E, Sarkar A, Shannon N, Stolte-Dijkstra I, Verloes A, Wakeling E, Wilson L, Hennekam RC. Nicolaides-Baraitser syndrome: Delineation of the phenotype. *Am J Med Genet A*. 2009 Aug;149A(8):1628-40. doi: 10.1002/ajmg.a.32956.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/19606471>
- Sousa SB, Hennekam RC; Nicolaides-Baraitser Syndrome International Consortium. Phenotype and genotype in Nicolaides-Baraitser syndrome. *Am J Med Genet C Semin Med Genet*. 2014 Sep;166C(3):302-14. doi: 10.1002/ajmg.c.31409. Epub 2014 Aug 28. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/25169058>
- Van Houdt JK, Nowakowska BA, Sousa SB, van Schaik BD, Seuntjens E, Avonce N, Sifrim A, Abdul-Rahman OA, van den Boogaard MJ, Bottani A, Castori M, Cormier-Daire V, Deardorff MA, Filges I, Fryer A, Fryns JP, Gana S, Garavelli L, Gillesen-Kaesbach G, Hall BD, Horn D, Huylebroeck D, Klapacki J, Krajewska-Walasek M, Kuechler A, Lines MA, Maas S, Macdermot KD, McKee S, Magee A, de Man SA, Moreau Y, Morice-Picard F, Obersztyn E, Pilch J, Rosser E, Shannon N, Stolte-Dijkstra I, Van Dijck P, Vilain C, Vogels A, Wakeling E, Wiczorek D, Wilson L, Zuffardi O, van Kampen AH, Devriendt K, Hennekam R, Vermeesch JR. Heterozygous missense mutations in SMARCA2 cause Nicolaides-Baraitser syndrome. *Nat Genet*. 2012 Feb 26;44(4):445-9, S1. doi: 10.1038/ng.1105.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/22366787>

Reprinted from Genetics Home Reference:

<https://ghr.nlm.nih.gov/condition/nicolaides-baraitser-syndrome>

Reviewed: December 2015

Published: March 21, 2017

Lister Hill National Center for Biomedical Communications
U.S. National Library of Medicine
National Institutes of Health
Department of Health & Human Services